

Conservative vs. Surgical intervention in Diaphyseal Forearm Fractures in age group 1 to 15 years: A Prospective Study

Rahul Madhukar Salunkhe¹, Sachin Sonawane¹, Sumeet Kumar¹, Anirudh Kandari¹

Abstract

Background: Pediatric diaphyseal fractures of the radius and ulna, commonly referred to as both bone forearm fractures, are the third most common fracture in the pediatric population and account for 13-40% of all pediatric fractures.^{1,2} Over 40% of girls and over 50% of boys sustain at least one fracture during childhood and adolescence,⁽³⁾ with distal forearm fractures being among the most common, accounting for up to one-third of all pediatric fractures.⁽³⁾ Furthermore, distal forearm fractures in childhood are increasing in incidence,^(4,5) particularly among girls.⁽⁴⁾ Fracture rate peaks between ages 11 and 15 yr [5], corresponding to the period of maximum postnatal growth velocity. Thus, the role of a childhood distal forearm fracture, in particular, on fracture risk later in life, would have practical clinical applications.

Introduction

Diaphyseal fracture of both bones of forearm in mature bone is now treated by open reduction and internal fixation. But it is not true for immature bones. Fractures of forearm in children and adolescent are extremely common.^[6] Studies have demonstrated lower bone density or weaker bone structure in boys and girls with a distal forearm fracture compared with controls.⁽⁷⁾

Recently there has been an increased interest in determining which method provides superior results, but the optimal treatment remains controversial.⁶ The goal of this paper is to study the current treatment of paediatric both bone forearm fractures in younger children (ages 1-15), and offer useful treatment algorithms for these injuries.^[8] Recently, however, there has been a trend towards increased surgical management of these fractures in an effort to improve clinical outcomes.^[9]

Material and Methods

- Institute Scientific & Ethics Committee Clearance was obtained before the start of the study. It was a prospective study of 50 cases done between the period of July, 2015 to September, 2017 at Dr D. Y. Patil Medical College, Pimpri, Pune.
- 50 patients with 50 forearm fractures, conservatively or operatively treated were included. The study was approved by the institutional ethical committee.
- Isolated radial head or neck fractures, multiple trauma, neurovascular injuries, Open fractures of Type 2 and 3 of Gustillo Anderson Classification, failure to achieve close reduction after three to four attempts, refractures, pathological Fractures and fractures with compartment syndrome were excluded

- Patients were divided into two groups the first group of patients were those who were treated successfully conservatively by manipulation and casting, the other group of patients were those who required open reduction and internal fixation. Variables were collected about the patients and methods of treatment (conservative versus open reduction and internal fixation) to find which of the two methods is more beneficial.

- **Nonoperative**
 - o closed reduction and immobilization
 - o indications: most pediatric forearm fractures can be treated without surgery
 - o greenstick injuries, bayonet apposition if <10 years
 - o follow-up with weekly radiographs for first 3-4 weeks to monitor reduction,
 - o casting is done for 6-12 weeks in total
 - o short arm cast for distal 1/3 BBFA
 - o above elbow immobilization for any fracture proximal to distal 1/3
- **Operative**
 - o **Percutaneous**
 - o indications: unacceptable alignment following closed reduction
 - o i.e
 - o angulation >15°, rotation >45° in children <10y
 - o angulation >10°, rotation >30° in children >10y
 - o bayonet apposition in children older than 10 years
 - o both bone forearm fractures in children > 13
 - o highly displaced fractures
 - o open reduction and internal fixation
 - o indications
 - o unacceptable alignment following closed reduction
 - o open fractures
 - o refractures
 - o angulation >15° and rotation >45° in children <10y

¹Department of Orthopaedics, Dr. D. Y. Patil Medical College, Hospital and; Research Centre, Pune, Maharashtra, India

Address for correspondence:

Dr. Anirudh Kandari

Department of Orthopaedics, Dr. D. Y. Patil Medical College, Hospital and; Research Centre, Pune, Maharashtra, India

Email: sunnykandari@hotmail.com

Table 1: Treatment of pediatric fractures[10]

Method	Remarks
Functional	Redression (e.g., collar-and-cuff bandage), functional bandage (Desault, Gilchrist) with acceptable degree of deformity
Immobilization	Plaster cast, conventional (white) or synthetic (rigid or semirigid; if necessary, correction by cast wedging)
Adaptation osteosynthesis (Kirschner wires)	Metaphyseal fractures: additional plaster cast immobilization necessary (unstable)
ESIN (elastic stable intramedullary nailing)	In longitudinally stable (transverse) diaphyseal fractures (also greenstick fractures of forearm shaft; as intramedullary rod in proximal upper arm fractures)
Plate fixation	As an exception, in fractures close to joints in adolescents
Medullary or locking nail	In diaphyseal fractures in adolescents

Table 2: Table of recommended acceptable alignment parameters for both-bone pediatric forearm fracture.[11]

Source	Age, years	Angulation, degrees	Malrotation, degrees	Bayonette apposition /displacement
Price (2010)15	<8	<15 MS, DS; <10 PS	<30	100% displacement
Noonan (1998)16	<9	<15	<45	<1 cm short
Tarmuzi (2009)17	<10	<20		No limits
Qairul (2001)20	<12	<20		

angulation >10° and rotation >30° in children >10y
 bayonet apposition in children older than 10 years
 both bone forearm fractures in children > 13
 highly displaced fractures

remaining(10.25) had good functional outcome results while of those children that had been treated operatively (80%) had an excellent, (16%) had good and (4%) had fair functional outcome results, this is according to Price et al.(10)

(Those children that needed internal fixation; we treat them either: By intra-medullary nails, K-wires cases or by plate and screws cases. The patient and method of fixation were chosen randomly).

Discussion

With the exception of severe fracture comminution, most both bone forearm fractures that can be treated by plate fixation may also be treated with flexible nails through closed or open reduction techniques. Recently fracture fixation with flexible nails has gained popularity, with proponents arguing that nailing results in decreased surgical dissection and retention of biologic factors at the fracture site.12,13 Both titanium and stainless steel flexible nails are available.

Given the excellent remodeling potential with younger patients, certain studies have argued that even with 100% displacement of the radius and ulna closed reduction and casting is an excellent treatment choice for children 9 years old and younger.14,15. However, the exact amount of angulation, displacement, and rotation that is acceptable remains controversial in the literature. It is generally accepted that the closer the fracture is to the distal physis, the greater the potential for remodeling. Consequently more deformity can be accepted in the distal one third of the diaphysis versus the middle and proximal thirds.

Observation and Results

Level of fracture site: Of those children (3.88%) had fracture in the proximal 1/3, (50.58%) had fracture in the middle 1/3 and (45.63) had fracture in the lower 1/3 of forearm bones.

Franklin et al. defined successful treatment of pediatric forearm fractures should result in painless and complication-free outcomes with functional pronosupination.15 It has been shown that 15 to 20 degrees of angulation in middle third forearm fractures can lead to major loss of forearm rotation.16 However, the significance of this range of motion loss as it pertains to clinical outcome remains debatable. Functional outcomes are satisfactory for closed management if manipulation can maintain reduction within this range17,18 Tarmuzi et al. concluded that up to 1cm of shortening can be accepted for closed management.17 Failure of closed management is rare, with

The functional outcome result: For the functional outcome results, of those children that had been treated conservatively (89.74%) had an excellent and the

Table 3: Complications: The main complications are:

	Conservative	Operative	Total
1. no complication	35(70%)	9(18%)	44(88%)
2. oedema	2(4%)	1(2%)	3(6%)
3. superficial & pin tract infection	0	1(2%)	1(2%)
4. compartment syndrome	0	0	0

Table 4: Complications: The main complications are:

Outcome	Conservative	Operative	Total
Excellent	70(67.96%)	20(19.41%)	90(87.37%)
Good	8(7.76%)	4(3.88%)	12(11.65%)
Fair	0	1(0.97)	1(0.97)
Poor	0	0	0
Total	78(75.72%)	25(24.27%)	103.00

Table 5: Table 12 Duration of hospitalization: Were longer for those treated by operative method, those patients kept in the ward under observation and care, for 1-2 days longer than those treated by conservative method, duration of stay in hospital for those treated by conservative method was (1-2) days, while for those treated by operative method was (2-3) days.

After treatment in day	Conservative	Operative	Total
1 day	73(70.87%)		73(70.87%)
2 day	5(4.85%)	21(20.38%)	26(25.24%)
3 day		4(3.88%)	4(3.88%)
More			
Total	78(75.72%)	25(24.27%)	103.00

roughly 90% of injuries being amenable to closed management.¹⁹ Children under the age of 4 should be placed in an above-elbow cast for any forearm fractures as short arm casts may slip.^{20,21} Post reduction, patients should be followed weekly for the first two to three weeks to ensure reduction is maintained. Holmes et al. found that if loss of reduction occurs, wedging the cast may restore alignment, but re-reduction or operative intervention may be required.²¹ The complications associated with cast immobilization include disuse osteopenia, muscle atrophy, skin breakdown, and elbow stiffness.^{22,23} Loss of reduction is the most common complication in pediatric forearm fractures, with rates between 10 and 60%.²²

Nonoperative management continues to be a very common, safe, and successful treatment option in pediatric forearm fractures. For those fractures that fail or are not amenable to conservative management however, surgical stabilization may need to be considered.

Different from other long bone of the human body, radius and ulna possess an important function of rotation, which play a crucial role for a series of nimble movements of the upper limb. The classic concept considers the double bones of forearm as a "joint" that allows rotation of the radius around the ulna, rather than two simple "long bone" [24, 25]. Holmes et al. stated that compression plating maximizes the ability to obtain anatomic reduction and restore normal radial bow.²¹ Additionally, given the construct fixation strength, plate fixation permits early range of motion. Although exact indications are debatable, it is suggested that plate fixation is indicated in the setting of significant comminution or with late loss of reduction after conservative management, as callous can prevent passage of intramedullary fixation.²¹ In contrast to adult fixation, smaller plate size and fewer screws can be used in children.^{21,30} Generally screw diameters are 2.7 mm or 3.5 mm, and 1/3 tubular plates and may be considered adequate.²⁶ Fixation through 4 cortices should be obtained proximal and distal to the fracture site, and the plate should not be wider than the bone.^{26,27}

.Complications of plate fixation include damage to surrounding structures, nonunion/malunion, and synostosis.²⁴ The rate of synostosis has been noted to have

an increased incidence if only one incision is used.²⁴ The potential for nerve damage is also present, more commonly secondary to ulnar fixation.³⁶

Other important factor in plating is implant removal, there was a 7.3% risk of an implant-related fracture in the follow up period, which all occurred within the first 3 years

With the exception of severe fracture comminution, most both bone forearm fractures that can be treated by plate fixation may also be treated with flexible nails through closed or open reduction techniques. Recently fracture fixation with flexible nails has gained popularity, with proponents arguing that nailing results in decreased surgical dissection and retention of biologic factors at the fracture site.²⁸ Both titanium and stainless steel flexible nails are available. In the clinical setting, titanium (Ti 6Al14V) is being used more often than stainless steel in most circumstances because of the elastic properties which allow for improved insertion and rotation while still providing adequate fracture stabilization.²⁹ Kang et al. evaluated 90 children treated with intramedullary nailing and reported good results and patient outcomes.³⁰ Complications secondary to intramedullary fixation include infection at the site of implantation, skin irritation, refracture after removal, implant failure, nerve/tendon injury, decreased range of motion, and compartment syndrome.

Conclusion

Conservative management is still the first line of treatment for pediatric forearm fractures especially in children less than 10 years old. Presently if operative intervention is required, both plate fixation and flexible nailing are acceptable treatment options. However, based on analysis of the available literature, it is unclear whether flexible nails or open reduction and internal fixation with plates should be recommended as a superior technique. Adequate understanding of the subtleties of either technique is necessary to ensure optimal outcomes, including the limitations of each technique and possible complications. In general, severe comminution and bone loss should be considered as indications for plate fixation, while intramedullary nailing offers better cosmesis, and decreased soft tissue disruption.

References

1. Cheng JC, Ng BK, Ying SY, Lam PK. A 10-year study of the changes in the pattern and treatment of 6,493 fractures. *J Pediatr Orthop.* 1999;19:344-50 [PubMed]
2. Jones K, Weiner DS. The management of forearm fractures in children: a plea for conservatism. *J Pediatr Orthop.* 1999;19:811-5 [PubMed]
3. Hedström EM, Svensson O, Bergström U, Michno P. Epidemiology of fractures in children and adolescents. *Acta Orthopaedica.* 2010;81 (1):148-53
4. Khosla S, Melton LJ, 3rd, Dekutoski MB, Achenbach SJ, Oberg AL, Riggs BL. Incidence of childhood distal forearm fractures over 30 years: a population-based study. *JAMA.* 2003;290(11):1479-85.
5. Mayranpaa MK, Makitie O, Kallio PE. Decreasing incidence and changing pattern of childhood fractures: a population-based study. [Erratum appears in *J Bone Miner Res.* 2011 Feb;26(2):439] *J Bone Miner Res.* 2009;25(12):2752-9
6. Campbell's operative orthopaedics: C.V. Mosbo Co. 7th Latest Edition, Vol.1, 2, 3, 4.
7. Kalkwarf HJ, Laor T, Bean JA. Fracture risk in children with a forearm injury is associated with volumetric bone density and cortical area (by peripheral QCT) and areal bone density (by DXA) *Osteoporos Int.* 2011;22(2):607-16.
8. Abraham A, Kumar S, Chaudhry S, Ibrahim T. Surgical interventions for diaphyseal fractures of the radius and ulna in children. *Cochrane Database Syst Rev* 2011:CD007907.
9. Flynn JM, Jones KJ, Garner MR, Goebel J. Eleven years experience in the operative management of pediatric forearm fractures. *J Pediatr Orthop.* 2010;30:313-9
10. *Dtsch Arztebl Int.* 2010 Dec; 107(51-52): 903-910.
11. *Orthop Rev (Pavia).* 2014 Apr 22; 6(2): 5325.
12. Sinikumpu JJ, Pokka T, Serlo W. The changing pattern of pediatric both-bone forearm shaft fractures among 86,000 children from 1997 to 2009. *Eur J Pediatr Surg.* 2013;23:289-96
13. Prevot J. [Stable elastic nailing]. *Z Unfallchir Versicherungsmed Berufskr.* 1989;82:252-60 [Article in French].
14. Zions LE, Zalavras CG, Gerhardt MB. Closed treatment of displaced diaphyseal both-bone forearm fractures in older children and adolescents. *J Pediatr Orthop.* 2005;25:507-12
15. Franklin CC, Robinson J, Noonan K, Flynn JM. Evidence-based medicine: management of pediatric forearm fractures. *J Pediatr Orthop.* 2012;32:S131-4
16. Shah AS, Lesniak BP, Wolter TD, et al. Stabilization of adolescent both-bone forearm fractures: a comparison of intramedullary nailing versus open reduction and internal fixation. *J Orthop Trauma.* 2010;24:440-7
17. Tarmuzi NA, Abdullah S, Osman Z, Das S. Paediatric forearm fractures: functional outcome of conservative treatment. *Bratisl Lek Listy.* 2009;110:563-8
18. Qairul IH, Kareem BA, Tan AB, Harwant S. Early remodeling in children's forearm fractures. *Med J Malaysia.* 2001;56:34-7
19. Smith VA, Goodman HJ, Strongwater A, Smith B. Treatment of pediatric both-bone forearm fractures: a comparison of operative techniques. *J Pediatr Orthop.* 2005;25:309-13
20. Do TT, Strub WM, Foad SL, et al. Reduction versus remodeling in pediatric distal forearm fractures: a preliminary cost analysis. *J Pediatr Orthop B.* 2003;12:109-15
21. Holmes JH, Wiebe DJ, Tataria M, et al. The failure of nonoperative management in pediatric solid organ injury: a multi-institutional experience. *J Trauma.* 2005;59:1309-13
22. Sinikumpu JJ, Pokka T, Serlo W. The changing pattern of pediatric both-bone forearm shaft fractures among 86,000 children from 1997 to 2009. *Eur J Pediatr Surg.* 2013;23:289-96
23. Ceroni D, Martin X, Delhumeau-Cartier C, et al. Is bone mineral mass truly decreased in teenagers with a first episode of forearm fracture? A prospective longitudinal study. *J Pediatr Orthop.* 2012;32:579-86
24. Jupiter J, Kellam J. Diaphyseal fractures of the forearm. In: Jupiter J, Browner B, editors. *SkellMN versus plates for adult BBFF 11370 Int J Clin Exp Med* 2017;10(8):11360-11370 etal *Trauma*, 4th edition. Philadelphia, PA: Saunders Elsevier; 2008. pp. 1459-1502.
25. Tarr RR, Garfinkel AI, Sarmiento A. The effects of angular and rotational deformities of both bones of the Forearm. An in vitro study. *J Bone Joint Surg Am* 1984; 66: 65-70
26. Wyrsh B, Mencio GA, Green NE. Open reduction and internal fixation of pediatric forearm fractures. *J Pediatr Orthop.* 1996;16:644-50
27. Bhaskar AR, Roberts JA. Treatment of unstable fractures of the forearm in children. Is plating of a single bone adequate? *J Bone Joint Surg Br.* 2001;83:253-8
28. Prevot J. [Stable elastic nailing]. *Z Unfallchir Versicherungsmed Berufskr.* 1989;82:252-60 [Article in French].
29. Lascombes P, Haumont T, Journeau P. Use and abuse of flexible intramedullary nailing in children and adolescents. *J Pediatr Orthop.* 2006;26:827-34
30. Kang SN, Mangwani J, Ramachandran M, et al. Elastic intramedullary nailing of paediatric fractures of the forearm: a decade of experience in a teaching hospital in the United Kingdom. *J Bone Joint Surg Br.* 2011;93:262-5

Conflict of Interest: NIL
Source of Support: NIL

How to Cite this Article

Salunkhe RA, Sonawane S, Kumar S, Kandari A. Conservative vs. Surgical intervention in Diaphyseal Forearm Fractures in age group 1 to 15 years: A Prospective Study. *Journal of Trauma and Orthopaedic Surgery* April-June 2018;13(2): 19-22